



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

the position of DH or BF . The equations of DH are $x=\frac{1}{2}\sqrt{2}$, $y=-\sqrt{2}z$, and the equations of BF are $x=-\frac{1}{2}\sqrt{2}$, $y=-\sqrt{2}z$. In either case

$x^2=\frac{1}{2}$ and $y^2=2z^2$ and $x^2+y^2=2z^2+\frac{a^2}{2}$ which is the equation

of the surface generated by the gauche hexagon $EHDCBF$.

This surface could also be generated by the hyperbola $x^2=2z^2+\frac{1}{2}$. Hence the volume of the hyperboloid of one


nappe generated $=\int \pi x^2 dz$, the upper limit being $\frac{1}{2}\sqrt{3}a$ and

the lower limit $-\frac{1}{2}\sqrt{3}a$. This integral is $\frac{5}{2}\pi\sqrt{3}a^3$.

The lines AB , AE , and AD generate a cone, radius $=\frac{1}{2}\sqrt{6}a$, altitude $=\frac{1}{2}\sqrt{3}a$, volume $=\frac{5}{2}\pi\sqrt{3}a^3$.

The lines GF , GH , and GC generate another cone of the same size.

The sum of the volumes of the three solids $=\frac{1}{2}\pi\sqrt{3}a^3=1.8138a^3$.



[NOTE.—This solution by the Proposer is fuller than that given in the November number, and is published because several of our contributors failed to comprehend the abbreviated solution previously published. Prof. Whitaker asserts that the solution by Dr. Zerr in the September-October number is incorrect, while the latter says he does not as yet see Prof. Whitaker's hyperboloid. The above seems to be correct, but we shall be glad to have the criticisms of other contributors.—EDITOR.]

43. Proposed by Professor C. E. WHITE, A. M., Trafalgar, Indiana.

Prove that $\int_0^1 \frac{x^{a-1} + x^{-a}}{1+x} \frac{dx}{x} = \log\left(\tan \frac{a\pi}{2}\right)$, when $a > 0$ and < 1 .

[Williamson's *Integral Calculus*, p. 154.]

Comment by C. W. M. BLACK, A. M., Professor of Mathematics in Wesleyan Academy, Wilbraham, Massachusetts.

There seems to be an error in No. 43, as I find the following in my copy of *Williamson* :


$$\int_0^1 \frac{x^{a-1} + x^{-a}}{1+x} \frac{dx}{\log x},$$

which gives the required result.

[In *Williamson's Integral Calculus*, edition of 1891, the problem is given as published, but the mistake has doubtless been corrected in the later edition.—EDITOR.]

44. Proposed by DE VOLSON WOOD, C. E., Professor of Mechanical and Electrical Engineering in Stevens Institute of Technology, Hoboken, New Jersey.

Find the equation of a curve in which $\rho = f(\theta)$, in which ρ is equal to BC , an intercept of any secant drawn from the corner E of the rectangle $AEDB$, and prolonged to cut AB prolonged in C . Let equal increments of θ be proportional to the equal increments of DB as divided by the secant EF , θ being zero when EC coincides with ED , and $\theta = 2\pi$ when EF passes through B . Determine the asymptotes.



I. Solution by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Irving College, Mechanicsburg, Pennsylvania; G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Applied Science, Texarkana College, Texarkana, Arkansas-Texas; and ALFRED HUME, C. E., D. Sc., Professor of Mathematics, University of Mississippi, University P. O., Mississippi.

Referring to the diagram given by the Proposer of this problem, July-Au-

gust (1895) MONTHLY, we have from the similar triangles FBC and FDE the following proportions: $BC : DE :: BF : DF$, or $\rho : b :: 2\pi - \theta : \theta$.

$\therefore (\rho + b)\theta = 2\pi b$ (1), which is the polar equation of *The Thistle of Scotland*, adopting the suggestion of Prof. MacCord.

Since $\rho^2(d\theta/d\rho) = -[(2\pi - \theta)^2/2\pi]b$, there is a *rectilinear asymptote* parallel to the initial line and at a distance $2\pi b$ above it. Making $\theta = \infty$, we have from (1) the equation $\rho = -b$; and this equation characterizes an *asymptotic circle* of radius b , or a *circular asymptote* of same radius, of the curve.

NOTE.—The derivation of (1) can be affected in, at least, *three* different ways; and, according to the conditions of the problem, (1) may also be written

$$(\rho + b)\theta = ab$$
 (2).

II. Solution by WILLIAM SYMMONDS, A. M., Professor of Mathematics and Astronomy, Pacific College, Santa Rosa, P. O. Sebastopol, California.

$$\text{From figure given } \frac{BC}{ED} = \frac{BF}{DF} = \frac{BD - DF}{DF} = \frac{BD}{DF} - 1,$$

$$\text{or } \frac{\rho}{b} = \frac{2\pi}{\theta} - 1; \quad \rho = \frac{2b\pi}{\theta} - b, \text{ the equation of the curve.}$$

When $\theta = 0$, $\rho = \infty$, and subtangent $= -2b\pi$.

The curve has, therefore, an asymptote parallel to OX at a distance above it, $2b\pi$, the circumference of a circle with radius AB .

The curve is concave toward the pole and intersects the axis perpendicularly and at a distance b to the left of the pole.

Elaborately solved by O. W. ANTHONY, and C. W. M. BLACK.

ERRATA.—On page 363, of last issue, line 4. omit $\sqrt{3}$ in the numerator of the second term; line 9, in the numerator, for “ $(a^2 + x^2)$ ” read $(a^2 - x^2)$; line 11, in the denominator of the second term, for “4” read 4^2 ; line 14, for “+” read $=$, before the last expression; page 364, line 15, for “of” read *to*; line 17, insert comma after “length”; line 17, for “ $2n$ ” read 2π ; line 18, for “ π_2 ” read π^2 ; on same page, problem No. “43” should be No. 42; page 365, line 1, for “ z^{n-2} ” read z^{n-1} ; and in line 2, of solution III., for “ $n^2 + y^2$ ” in the exponent, read $x^2 + y^2$.

PROBLEMS.

51. Proposed by F. P. MATZ, D. Sc., Ph. D., Professor of Mathematics and Astronomy in Irving College, Mechanicsburg, Pennsylvania.

Find the maximum ellipsoid that can be cut out of a given right conic frustrum.

52. Proposed by O. W. ANTHONY, M. Sc., Professor of Mathematics in New Windsor College, New Windsor, Maryland.

There are two lights of intensities m and n . Where must a target, whose surface is parallel to the line joining the two lights, be set up in order that it shall receive the maximum illumination per unit of area.